

TOYOPEARL Butyl-600M

TOYOPEARL Butyl-650C

TOYOPEARL Butyl-650M

TOYOPEARL Butyl-650S

TOYOPEARL Ether-650M

TOYOPEARL Ether-650S

TSKgel Ether-5PW (20)

TSKgel Ether-5PW (30)

TOYOPEARL Hexyl-650C

TOYOPEARL Phenyl-600M

TOYOPEARL Phenyl-650C

TOYOPEARL Phenyl-650M

TOYOPEARL Phenyl-650S

TSKgel Phenyl-5PW (20)

TSKgel Phenyl-5PW (30)

TOYOPEARL PPG-600M

TOYOPEARL SuperButyl-550C

The role of Hydrophobic Interaction Chromatography in Process Purification

Hydrophobic interaction chromatography (HIC) is a powerful tool for the process purification of biomolecules. The technique utilizes the accessible hydrophobic regions located on protein surfaces and their interactions with a weakly hydrophobic stationary phase. HIC is an excellent complement to ion exchange and size exclusion chromatography particularly when protein isoforms exist or when feedstock impurities are of similar isoelectric point or molar mass. The selectivity differences exploited by HIC can also be used after affinity separations in which closely related proteins with similar recognition sites are not distinguishable by the affinity ligand.


Proteins and other molecules with hydrophobic surfaces are attracted to the hydrophobic ligands of HIC resins. Proteins are bound to the resin by employing an aqueous high salt mobile phase. The salt conditions contribute to a lyotropic effect which allows the proteins to bind to the lower surface coverage of a hydrophobic ligand. Proteins are eluted by the simple technique of decreasing the salt concentration. Most therapeutic targets are eluted in a low salt or a no salt buffer.

During elution, the energy of interaction for a HIC step is less than that of a reversed phase chromatography (RPC) step. One means of gauging the relative binding energy between the two techniques is to measure the surface tension of the two sets of binding and elution conditions. **Figure 1** provides a comparison of the surface tension generated by HIC and RPC elution systems.¹ Since HIC separates under milder eluting conditions, biological activity is typically retained.

TOYOPEARL Hydrophobic Interaction Chromatography Resins

TOYOPEARL HIC resins are functionalized versions of the TOYOPEARL HW size exclusion resins and are therefore based on hydroxylated polymethacrylic polymer beads. Tosoh Bioscience offers five HIC ligands featuring different degrees of hydrophobicity and selectivity. **Table 1** lists the properties of these TOYOPEARL HIC resins. The hydrophobicity of TOYOPEARL HIC resins increases through the ligand series: ether, PPG (polypropylene glycol), phenyl, butyl, and hexyl (**Figure 2**).


Figure 1: The surface tension of aqueous solutions used in HIC and RPC



Mode	Gradient (Typical)	Δ Surface Tension (erg/cm ²)
HIC	1.8 to 0 mol/L (NH ₄) ₂ SO ₄ / aqueous buffer	4
RPC	10 to 50% ACN/ 0.1%TFA	23

¹C. Horvath et. al., Separation Processes in Biotechnology, Volume 9; Asenjo, J. ed.; Marcel Dekker, Inc.: New York, 1990, p 447.

Figure 2: Available HIC ligands







Table 1: Properties of TOYOPEARL HIC resins

TOYOPEARL resin	Hydrophobicity	Base bead	Pore size (nm)	Bead diameter (µm)	Ligand type	DBC (g/L)	Pressure rating
Ether-650S	+	HW-65	100	20 - 50	Ether	10-30	0.3 MPa
Ether-650M	+	HW-65	100	40 - 90	Ether	10-30	0.3 MPa
PPG-600M	++	HW-60	75	40 - 90	Polypropylene glycol	45 - 55	0.3 MPa
Phenyl-600M	+++	HW-60	75	40 - 90	Phenyl	45 - 65	0.3 MPa
Phenyl-650S	+++	HW-65	100	20 - 50	Phenyl	30 - 50	0.3 MPa
Phenyl-650M	+++	HW-65	100	40 - 90	Phenyl	30 - 50	0.3 MPa
Phenyl-650C	+++	HW-65	100	50 - 150	Phenyl	30 - 50	0.3 MPa
Butyl-650S	++++	HW-65	100	20 - 50	Butyl	30 - 50	0.3 MPa
Butyl-650M	++++	HW-65	100	40 - 90	Butyl	30 - 50	0.3 MPa
Butyl-650C	++++	HW-65	100	50 - 150	Butyl	30 - 50	0.3 MPa
Butyl-600M	++++	HW-60	75	40 - 90	Butyl	40 - 60	0.3 MPa
SuperButyl-550C	++++	HW-55	50	50 - 150	Butyl	52 - 70	0.3 MPa
Hexyl-650C	+++++	HW-65	100	50 - 150	Hexyl	30 - 50	0.3 MPa

Three HIC ligands are available in the TOYOPEARL -600 resin format: PPG, phenyl, and butyl. The selectivities of TOYOPEARL Butyl-600M, TOYOPEARL PPG-600M and the TOYOPEARL Phenyl-600M resins are shown in Figure 3. Available in the TOYOPEARL -650 series are the following four HIC ligands: hexyl, butyl, phenyl, and ether. The remaining ligand available in the TOYOPEARL HIC resin line is SuperButyl-550.

Figure 3: Comparison of TOYOPEARL -600M resins



Resins:
A. TOYOPEARL PPG-600M
B. TOYOPEARL Phenyl-600M
C. TOYOPEARL Butyl-600M

Column size: 7.5 mm ID × 7.5 cm

Mobile phase: Buffer A: 1.8 mol/L (NH₄)₂SO₄ + 0.1 mol/L sodium phosphate, pH 7.0
 Buffer B: 0.1 mol/L sodium phosphate, pH 7.0

Gradient: 60 min linear gradient from buffer A to B

Flow rate: 136 cm/hr (1.0 mL/min)

Detection: UV @ 280 nm


Temperature: ambient

Samples: 1 g/L of: 1. ribonuclease A 2. lysozyme
 3. α-chymotrypsinogen

Load volume: 100 µL

A comparison of the dynamic binding capacities (DBCs) of the TOYOPEARL -600 resins with TOYOPEARL Phenyl-650M is shown in Figure 4. Figure 5 compares the selectivities of the TOYOPEARL Phenyl-600M and TOYOPEARL Phenyl-650M resins with an agarose based phenyl resin. The narrower pore diameter of TOYOPEARL SuperButyl-550C resin (based on the 50 nm pore diameter TOYOPEARL HW-55 resin) is recommended for the analysis of smaller molecules such as lysozyme (1.2×10^4 Da). A comparison of the DBC of TOYOPEARL SuperButyl-550C resin with other TOYOPEARL HIC resins is shown in Figures 6 and 7.

Figure 4: Breakthrough curves of polyclonal IgG on various HIC resins




Resins:
A. TOYOPEARL PPG-600M
B. TOYOPEARL Phenyl-600M
C. TOYOPEARL Butyl-600M
D. Competitor Phenyl Agarose
E. TOYOPEARL Phenyl-600M

Column size: 7.8 mm ID \times 20 cm
 Mobile phase: 0.1 mol/L sodium phosphate, pH 7.0 + 0.8 mol/L $(\text{NH}_4)_2\text{SO}_4$
 Flow rate: 300 cm/hr (2.4 mL/min)
 Detection: UV @ 280 nm
 Temperature: 25 °C
 Samples: 1.0 g/L polyclonal IgG

DBC was calculated at 10% breakthrough

Figure 5: Selectivity comparison of phenyl-type resins



Resins:
A. TOYOPEARL Phenyl-600M
B. TOYOPEARL Phenyl-650M
C. Competitor Phenyl Agarose

Column size: 7.5 mm ID \times 7.5 cm
 Mobile phase: Buffer A: 1.8 mol/L $(\text{NH}_4)_2\text{SO}_4$ + 0.1 mol/L sodium phosphate, pH 7.0
 Buffer B: 0.1 mol/L sodium phosphate, pH 7.0
 Gradient: 60 min linear gradient from buffer A to B
 Flow rate: 136 cm/hr (1.0 mL/min)
 Detection: UV @ 280 nm
 Temperature: ambient
 Sample: 1.0 g/L of: 1. ribonuclease A 2. lysozyme
 3. α -chymotrypsinogen
 Load volume: 100 μ L





Figure 6: Typical dynamic binding capacities for lysozyme




Resins:	binding capacity (g/L) (10% breakthrough)
---------	--

A. TOYOPEARL Ether-650M	12.5
B. TOYOPEARL Phenyl-650M	27.5
C. TOYOPEARL Butyl-650M	32.2
D. TOYOPEARL Hexyl-650C	33.2
E. TOYOPEARL PPG-600M	44.2
F. TOYOPEARL SuperButyl-550C	54.3

Column size: 7.8 mm ID × 20 cm
 Mobile phase: 1.8 mol/L sodium sulfate + 0.1 mol/L phosphate, pH 7.0
 Flow rate: 100 cm/hr (0.8 mL/min)
 Detection: UV @ 280 nm
 Temperature: ambient
 Sample: 1 g/L lysozyme
 Sample load: as indicated in figure

Figure 7: TOYOPEARL Phenyl-600M breakthrough curve (lysozyme)



Resins:	binding capacity (g/L) (10% breakthrough)
---------	--

A. TOYOPEARL Phenyl-600M	58
B. TOYOPEARL Phenyl-650M	27

Column size: 7.8 mm ID × 20 cm
 Mobile phase: 1.8 mol/L (NH₄)₂SO₄ + 0.1 mol/L phosphate, pH 7.0
 Flow rate: 300 cm/hr (2.4 mL/min)
 Detection: UV @ 280 nm
 Temperature: ambient
 Sample: 1 g/L lysozyme
 Sample load: as indicated in figure

The larger pore TOYOPEARL products such as TOYOPEARL Butyl-650 and TOYOPEARL Phenyl-650 resins are very useful for protein aggregate separation and removal. In addition, Tosoh Bioscience HIC resins are very effective in separating misfolded proteins from the native protein form. Because misfolded proteins will generally be more hydrophobic than the native protein, TOYOPEARL Butyl-650M resin is used frequently for the removal of misfolded proteins. In many cases, flow-through chromatography can be accomplished under eluent conditions binding the misfolded protein while allowing the native target protein to flow through the column.

Hydrophobic interaction is a very useful technique for the purification of monoclonal antibodies (mAbs), with their diverse hydrophobic nature. The range of HIC ligands of varying hydrophobicity available from Tosoh Bioscience (Figure 2) gives chromatographic developers a range of options for finding the right ligand for their target molecule.

TSKgel Hydrophobic Interaction Chromatography Resins

The same ether and phenyl ligands that are used for the TOYOPEARL resins are also available within the TSKgel HIC resin product line. Properties of TSKgel HIC resins are listed in [Table 2](#). The TSKgel HIC resins use the same methacrylic polymer chemistry as the TOYOPEARL resins ([Table 3](#)) but have a higher degree of crosslinking, making for a more rigid bead. This is necessitated by the higher pressures generated when using smaller particles for chromatography. Greater crosslinking decreases the number of sites available for ligand attachment and thus a TSKgel resin will have a lower dynamic binding capacity than the corresponding TOYOPEARL resin. The polymeric structure of these products also makes them resistant to a wide range of pH conditions and mobile phase ionic strengths. In addition, the hydroxylated surface of the base bead reduces non-specific binding of proteins.

Table 2: Properties of TSKgel HIC resins


TSKgel resin	Hydrophobicity	Base bead	Pore size (nm)	Bead diameter (µm)	Ligand type	DBC (g/L)	Pressure rating
Ether-5PW (20)	+	PW5000	100	15 - 25	Ether	10 - 30	2.0 MPa
Ether-5PW (30)	+	PW5000	100	20 - 40	Ether	10 - 30	2.0 MPa
Phenyl-5PW (20)	++	PW5000	100	15 - 25	Phenyl	10 - 30	2.0 MPa
Phenyl-5PW (30)	++	PW5000	100	20 - 40	Phenyl	10 - 30	2.0 MPa

Table 3: Methacrylic base beads available for HIC

Pore size (nm)	5	12.5	40-50	75	100	>100	>170
Resin							
TOYOPEARL HW-type:	40	50	55	60	65	75	80
TSKgel PW-type:	G1000	G2000	G4000		G5000	G6000	

← Increasing pore surface area

TOYOPEARL HIC resins are chemically stable from pH 1-13. This allows a constant packing volume over a wide range of salt concentrations and cleaning in place (CIP) with acid or base. Also, these resins can be run at elevated temperatures (4-60 °C) and are autoclavable at 121 °C.



Because TOYOPEARL and TSKgel HIC resins have the same backbone polymer chemistry, the selectivity for proteins and impurities will be unchanged. **Table 4** shows the ligands and particle sizes available for TOYOPEARL and TSKgel HIC resins and is arranged in increasing levels of resolution by bead size (i.e. low, medium, and high resolution). The semi-rigid polymeric backbone of TOYOPEARL and TSKgel HIC resins permits high flow rates for maximum throughput and productivity. TOYOPEARL HIC resins may be operated at pressures up to 0.3 MPa and TSKgel -5PW HIC resins may be operated up to 2.0 MPa. The pressure-flow characteristics for each particle size grade of TOYOPEARL Phenyl-650 resins are shown in **Figure 8**.


Resolution increases with decreasing particle size. Resin particle size is proportional to HETP and inversely proportional to the column efficiency and resolution of two peaks. TOYOPEARL HIC resins are available in three particle sizes, though not all ligands are available in each grade:

- S-grade = 35 μm (Superfine)
- M-grade = 65 μm (Fine)
- C-grade = 100 μm (Coarse)

Some processes, such as the purification of antibody-drug conjugates, require resins that are capable of higher resolution separations. For these separations, smaller diameter TOYOPEARL S-grade or TSKgel resins are preferred. TSKgel HIC resins are currently available in two ligands and two bead sizes:

- TSKgel Ether-5PW (30) = 30 μm
- TSKgel Ether-5PW (20) = 20 μm
- TSKgel Phenyl-5PW (30) = 30 μm
- TSKgel Phenyl-5PW (20) = 20 μm

Figure 8: Pressure-flow curve for TOYOPEARL Phenyl-650 resins of various particle sizes



Resins:
A. TOYOPEARL Phenyl-650S
B. TOYOPEARL Phenyl-650M
C. TOYOPEARL Phenyl-650C

Column size: 25 mm ID \times 25 cm
 Mobile phase: 2.0 mol/L $(\text{NH}_4)_2\text{SO}_4$
 Flow rate: as indicated in figure

Table 4: Resolution of TOYOPEARL and TSKgel HIC resins

Resolution	Bead diameter (μm)	Pore size (nm)	HIC resin
Low	100	50	TOYOPEARL SuperButyl-550C
		100	TOYOPEARL Hexyl-650C
		100	TOYOPEARL Butyl-650C
		100	TOYOPEARL Phenyl-650C
Medium	65	75	TOYOPEARL Butyl-600M
		75	TOYOPEARL Phenyl-600M
		75	TOYOPEARL PPG-600M
	65	100	TOYOPEARL Butyl-650M
		100	TOYOPEARL Phenyl-650M
		100	TOYOPEARL Ether-650M
High	35	100	TOYOPEARL Butyl-650S
		100	TOYOPEARL Phenyl-650S
		100	TOYOPEARL Ether-650S
	30	100	TSKgel Phenyl-5PW (30)
		100	TSKgel Ether-5PW (30)
	20	100	TSKgel Phenyl-5PW (20)
100		TSKgel Ether-5PW (20)	

Parameters to Consider when Using Tosoh Bioscience HIC Resins

Coordinating the hydrophobicity of the therapeutic target to the resin hydrophobicity is critical for the best overall purification performance. Too hydrophobic a resin for a given protein can result in its irreversible binding to the resin or a loss of biological activity. **Tables 5 and 6** show typical mass recovery and biological activity recovery data for TOYOPEARL HIC resins.

Table 5: High mass recovery (%) of proteins

Protein	TOYOPEARL resin		
	Ether-650M	Phenyl-650M	Butyl-650M
bovine serum albumin	84	62	76*
α -chymotrypsinogen	96	88*	90
cytochrome c	—	81*	87*
IgG	91	—	—
α -lactalbumin	90	—	—
lysozyme	94	92	85
ovalbumin	83	88	73
ribonuclease A	—	72*	82*

Procedure: A 200 mL sample containing 200 mg of protein was loaded onto a 7.5 mm ID \times 7.5 cm column and eluted with a 60 minute gradient of 1.8 mol/L (*1.5 mol/L) to 0.0 mol/L ammonium sulfate in 0.1 mol/L sodium phosphate, pH 7.0. The mass recovery was determined spectrophotometrically at UV 280 nm and 25 °C.

Table 6: Recovery of enzymatic activity of proteins

TOYOPEARL resin	Protein	% Activity recovery
Phenyl-650	phytochrome	79
Butyl-650	halophilic protease	85
Butyl-650	poly (3-hydroxybutyrate) depolymerase	88
Butyl-650	aculeacin-A acylase	82
Butyl-650	opine dehydrogenase	81


An optimum HIC process step will balance high dynamic binding capacity, adequate selectivity, good mass recovery and retention of biological activity. The wide range of selectivities for TOYOPEARL and TSKgel resins enables a developer to optimize protein separations at the extremes of the hydrophobic spectrum. The more hydrophobic ligands on TOYOPEARL Hexyl-type and TOYOPEARL Butyl-type resins are used to separate hydrophilic proteins. These two resins should also be considered for separations requiring a low salt environment.

TOYOPEARL and TSKgel Ether resins are used for the purification of very hydrophobic targets such as certain monoclonal antibodies and membrane proteins. These proteins may bind irreversibly to other more hydrophobic resins.

TOYOPEARL PPG and TOYOPEARL and TSKgel Phenyl resins complement the other HIC ligands available in the HIC series and offer alternatives for moderately hydrophobic proteins.

In addition to the hydrophobicity of the ligand, the selectivity in HIC is influenced by the eluent salt type. **Figure 9** demonstrates the effect of salt type on the resolution factor of different protein pairs.² The Hofmeister lyotropic salt series shown in **Figure 10** ranks anions and cations by their ability to promote protein precipitation. Ions on the left are referred to as “lyotropic” while the ions on the right are called “chaotropic”. Lyotropic salts will precipitate or “salt out” proteins at high salt concentrations due to increased hydrophobic interaction, while chaotropic salts will promote protein denaturation at high salt concentrations. The Hofmeister lyotropic salt series indicates that the use of different salt systems may generate a variety of adsorption and desorption selectivities for each resin with a given protein. This feature of HIC provides an additional parameter for the optimization of a process step.

Figure 9: Influence of salt-type on resolution



Chromatography on a Toyopearl Butyl-substituted support

Resin: Toyopearl Butyl-650M
Column size: 4.1 mm ID \times 4 cm
Mobile phase: Buffer A: 20 mmol/L phosphate buffer in 1.0 mol/L indicated salt, pH 7.0
 Buffer B: buffer A with 1.0 mol/L indicated salt
Flow rate: 484 cm/hr (1 mL/min)
Detection: UV @ 280 nm

²Fausnaugh, J.; Kennedy, L.; Regnier, F. *J. Chromatography*, **1984**, 141, 317.




Figure 10: Hofmeister lyotropic salt series

for anions
 $\text{SO}_4^{2-} > \text{HPO}_4^{2-} > \text{CH}_3\text{COO}^- > \text{halide} > \text{NO}_3^- > \text{ClO}_4^- > \text{SCN}^-$

for cations
 $(\text{CH}_3)_4\text{N}^+ > \text{K}^+ > \text{Na}^+ > \text{Cs}^+ > \text{Li}^+ > \text{Mg}^{2+} > \text{Ca}^{2+} > \text{Ba}^{2+}$

Ammonium sulfate and sodium sulfate are the most commonly used salts in HIC. NaCl is often used as well.


HIC is commonly used as a polishing step in monoclonal antibody purification processes. HIC offers an orthogonal selectivity to ion exchange chromatography and can be an effective step for aggregate clearance and host cell protein reduction, however, this mode of chromatography suffers from the limitation of use of high concentrations of kosmotropic salts to achieve the desired separation. Ghose et al³ reports an unconventional way of operating HIC in the flowthrough (FT) mode with no kosmotropic salt in the mobile phase. TOYOPEARL Hexyl-650C was selected as the stationary phase and the pH of the mobile phase was modulated to achieve the required selectivity. Optimum pH conditions were chosen under which the antibody product of interest flowed through while impurities such as aggregates and host cell proteins bound to the column. The performance of the TOYOPEARL Hexyl-650C resin was comparable to that observed using conventional HIC conditions with high salt.

³Ghose, S.; Tao, Y.; Conley, L.; Cecchini, D. Purification of monoclonal antibodies by hydrophobic interaction chromatography under no-salt conditions. *mAbs*. 2013, 5, (5), 795-800.

Purification of Monoclonal Antibodies

For a very hydrophobic mAb, such as mouse anti-chicken lectin (14 kDa), the less hydrophobic TOYOPEARL Ether ligand works quite well. The purification of this mAb from ascites fluid (Figure 11) was performed with a 10 μm TSKgel Ether-5PW semi-preparative column. Identical selectivity for scale-up was found with corresponding 65 μm TOYOPEARL Ether-650M resin.

Figure 11: Purification of mAbs from ascites fluid



Resins: **A: TSKgel Ether-5PW (prepacked HPLC column)**
B: Toyopearl Ether-650M

Column size: 7.5 mm ID × 7.5 cm

Mobile phase: Buffer A: 1.5 mol/L $(\text{NH}_4)_2\text{SO}_4$ + 0.1 mol/L phosphate, pH 7.0
 Buffer B: 0.1 mol/L phosphate, pH 7.0

Gradient: 60 min linear gradient from buffer A to B

Flow rate: 136 cm/hr (1.0 mL/min)

Detection: UV @ 280 nm

Temperature: ambient


Sample: A: 1.5 mg/100 μL anti-chicken 14 kDa lectin
 B: 0.76 mg/50 μL diluted ascites fluid

Load volume: 150 μL

Plasmid DNA Purification

TOYOPEARL Hexyl-650C resin was used successfully for plasmid DNA purification by Cambrex, Baltimore, MD (US patent 6,953,686). The resin was shown to be the most effective among HIC resins for endotoxin removal with capacities exceeding 2 million EU/mL of resin. Additionally, RNA and protein impurities were effectively eliminated. TOYOPEARL Hexyl-650C was also effective in separating the supercoiled and open circular forms of plasmid DNA (Figure 12). Under certain binding conditions, the two forms are bound to the resin, and subsequently eluted with a simple gradient, resulting in two distinct peaks corresponding to the relaxed and supercoiled forms respectively.

Figure 12: Plasmid DNA separation




Resin: TOYOPEARL Hexyl-650C
Column size: 1.0 cm ID × 30 cm (23.6 mL)
Mobile phase: Buffer A. 3.0 mol/L ammonium sulfate, 10 mmol/L Tris, 1 mmol/L EDTA, pH 7.4
 Buffer B. 10 mmol/L Tris, 1 mmol/L EDTA, pH 7.4
Gradient: 3.0 mol/L ammonium sulfate to 1.0 mol/L ammonium sulfate (6 CV)
Flow rate: 153 cm/hr (2.0 mL/min)
Detection: UV @ 254 nm
Sample: Plasmid DNA in 3.0 mol/L ammonium sulfate


Purification of Glycoproteins

TOYOPEARL HIC resins can purify glycoproteins, which often bind irreversibly to saccharide-based chromatographic media. Figure 13 shows the purification of a large glycoprotein on TOYOPEARL Butyl-650S resin.

Figure 13: Large glycoprotein purified on TOYOPEARL Butyl-650S




Resin: Toyopearl Butyl-650S
Column size: 22 mm ID × 26 cm
Mobile phase: Buffer A: 40% saturated (NH₄)₂SO₄ + 50 mol/L phosphate, pH 7.0
 Buffer B: 50 mol/L phosphate, pH 7.0
Gradient: A: Load and wash in 100% buffer A
 B: 50% buffer A with 50% buffer B
 C: 100% buffer B
Detection: UV @ 280 nm
Sample: crude protein from sea hare *Aplysia kurodai*



Ultra Purification of Target Compound

Biopharmaceutical process development often requires a high performance step for ultra-purification of a target compound. To meet these needs, 20 and 30 μm TSKgel Phenyl-5PW and Ether-5PW are available. The selectivity of these packings is similar to the 10 μm TSKgel 5PW Phenyl-5PW and Ether-5PW analytical columns. Therefore methods can easily be transferred from analytical to preparative scale resins of the same chemistry using a seamless scale-up strategy. **Figure 14** shows the similar elution pattern on 10 μm and 30 μm TSKgel packings, along with 65 μm TOYOPEARL process-scale resin.

Figure 14: Seamless scale up



Resins: **A and B. TSKgel Phenyl-5PW**
C. TOYOPEARL Phenyl-650M

Column size: 7.5 mm ID \times 7.5 cm

Mobile phase: A and B: 60 min linear gradient from 1.8 mol/L to 0 mol/L $(\text{NH}_4)_2\text{SO}_4$ in 0.1 mol/L phosphate buffer

Gradient: C: 240 min linear gradient from 1.8 mol/L to 0 mol/L $(\text{NH}_4)_2\text{SO}_4$ in 0.1 mol/L phosphate buffer

Flow rate: A and B. 68 cm/hr (0.50 mL/min)
C. 272 cm/hr (2.0 mL/min)

Detection: UV @ 280 nm


Samples: 1. myoglobin
2. ribonuclease A
3. lysozyme
4. α -chymotrypsinogen

Load volume: 100 μL containing 1 g/L of each protein

Purification and Resolution of Pullulanase

The power of HIC is illustrated in a scheme in which pullulanase, an amylase-like enzyme responsible for hydrolysis of branched chain sugars, is purified and resolved into two closely related forms. Ion exchange and size exclusion chromatography effectively purified pullulanase. With TOYOPEARL Butyl-650S, however, two closely related proteins were resolved, based on differences in their surface hydrophobicity (Figure 15).

Figure 15: Separation of two active pullulanase forms




Resin: TOYOPEARL Butyl-650S
Column size: 18 mm ID × 16 cm
Mobile phase: isocratic elution, 120 mL (NH₄)₂SO₄, 25% saturation in 0.02 mol/L phosphate, pH 7.0, followed by a linear gradient, 224 mL (NH₄)₂SO₄, 25% to 0% saturation, in buffer
Flow rate: 12 cm/hr
Detection: UV @ 280 nm
Sample: protein from *Bacillus acidopullulyticus*
Sample load: 20 mg


Lipase Isozymes

Incorporation of HIC into a purification scheme has separated lipase isozymes that were not resolved by a previously reported method. After ion exchange and size exclusion chromatography, an additional step employing TOYOPEARL Butyl-650M, as shown in Figure 16, enabled the separation of two active lipase isozymes, L1 and L2, from an inactive impurity. Activity recovery was 93% for this step.

Figure 16: Separation of lipase isozymes from impurity



Resin: TOYOPEARL Butyl-650M
Column size: 34 mm ID × 29 cm
Mobile phase: linear gradient, 810 mL (NH₄)₂SO₄, 25% to 0% saturation in 0.01 mol/L acetate buffer, pH 5.6
Flow rate: 2 cm/hr
Detection: UV @ 254nm
Sample: lipase from *Geotrichum candidum*
Sample load: 375 mg




A selection of screening tools are available for TOYOPEARL and TSKgel HIC resins. See the Process Development Products section of this Product Guide for details.

Ordering Information

TOYOPEARL HIC resins:

Part #	Product description	Container size (mL)	Bead diameter (µm)	Typical lysozyme capacity (g/L)	
43151	TOYOPEARL Ether-650S	25	20 - 50	10 - 30	
16172	TOYOPEARL Ether-650S	100	20 - 50	10 - 30	
16174	TOYOPEARL Ether-650S	1,000	20 - 50	10 - 30	
16176	TOYOPEARL Ether-650S	5,000	20 - 50	10 - 30	
19805	TOYOPEARL Ether-650M	25	40 - 90	10 - 30	
16173	TOYOPEARL Ether-650M	100	40 - 90	10 - 30	
16175	TOYOPEARL Ether-650M	1,000	40 - 90	10 - 30	
16177	TOYOPEARL Ether-650M	5,000	40 - 90	10 - 30	
21301	TOYOPEARL PPG-600M	25	40 - 90	45 - 55	
21302	TOYOPEARL PPG-600M	100	40 - 90	45 - 55	
21303	TOYOPEARL PPG-600M	1,000	40 - 90	45 - 55	
21304	TOYOPEARL PPG-600M	5,000	40 - 90	45 - 55	
21305	TOYOPEARL PPG-600M	50,000	40 - 90	45 - 55	
21887	TOYOPEARL Phenyl-600M	25	40 - 90	45 - 65	
21888	TOYOPEARL Phenyl-600M	100	40 - 90	45 - 65	
21889	TOYOPEARL Phenyl-600M	1,000	40 - 90	45 - 65	
21890	TOYOPEARL Phenyl-600M	5,000	40 - 90	45 - 65	
20891	TOYOPEARL Phenyl-600M	50,000	40 - 90	45 - 65	
43152	TOYOPEARL Phenyl-650S	25	20 - 50	30 - 50	
14477	TOYOPEARL Phenyl-650S	100	20 - 50	30 - 50	
14784	TOYOPEARL Phenyl-650S	1,000	20 - 50	30 - 50	
14935	TOYOPEARL Phenyl-650S	5,000	20 - 50	30 - 50	
19818	TOYOPEARL Phenyl-650M	25	40 - 90	30 - 50	
14478	TOYOPEARL Phenyl-650M	100	40 - 90	30 - 50	
14783	TOYOPEARL Phenyl-650M	1,000	40 - 90	30 - 50	
14943	TOYOPEARL Phenyl-650M	5,000	40 - 90	30 - 50	
18364	TOYOPEARL Phenyl-650M	50,000	40 - 90	30 - 50	
43126	TOYOPEARL Phenyl-650C	25	50 - 150	30 - 50	
14479	TOYOPEARL Phenyl-650C	100	50 - 150	30 - 50	
14785	TOYOPEARL Phenyl-650C	1,000	50 - 150	30 - 50	
14944	TOYOPEARL Phenyl-650C	5,000	50 - 150	30 - 50	

Part #	Product description	Container size (mL)	Bead diameter (µm)	Typical lysozyme capacity (g/L)	
43153	TOYOPEARL Butyl-650S	25	20 - 50	30 - 50	
07476	TOYOPEARL Butyl-650S	100	20 - 50	30 - 50	
14701	TOYOPEARL Butyl-650S	1,000	20 - 50	30 - 50	
07975	TOYOPEARL Butyl-650S	5,000	20 - 50	30 - 50	
18826	TOYOPEARL Butyl-650S	50,000	20 - 50	30 - 50	
19802	TOYOPEARL Butyl-650M	25	40 - 90	30 - 50	
07477	TOYOPEARL Butyl-650M	100	40 - 90	30 - 50	
14702	TOYOPEARL Butyl-650M	1,000	40 - 90	30 - 50	
07976	TOYOPEARL Butyl-650M	5,000	40 - 90	30 - 50	
18355	TOYOPEARL Butyl-650M	50,000	40 - 90	30 - 50	
43127	TOYOPEARL Butyl-650C	25	50 - 150	30 - 50	
07478	TOYOPEARL Butyl-650C	100	50 - 150	30 - 50	
14703	TOYOPEARL Butyl-650C	1,000	50 - 150	30 - 50	
07977	TOYOPEARL Butyl-650C	5,000	50 - 150	30 - 50	
22826	TOYOPEARL Butyl-650C	50,000	50 - 150	30 - 50	
21448	TOYOPEARL Butyl-600M	25	40 - 90	40 - 60 (γ-globulin)	
21449	TOYOPEARL Butyl-600M	100	40 - 90	40 - 60 (γ-globulin)	
21450	TOYOPEARL Butyl-600M	1,000	40 - 90	40 - 60 (γ-globulin)	
21451	TOYOPEARL Butyl-600M	5,000	40 - 90	40 - 60 (γ-globulin)	
21452	TOYOPEARL Butyl-600M	50,000	40 - 90	40 - 60 (γ-globulin)	
19955	TOYOPEARL SuperButyl-550C	25	50 - 150	52 - 70	
19956	TOYOPEARL SuperButyl-550C	100	50 - 150	52 - 70	
19957	TOYOPEARL SuperButyl-550C	1,000	50 - 150	52 - 70	
19958	TOYOPEARL SuperButyl-550C	5,000	50 - 150	52 - 70	
19959	TOYOPEARL SuperButyl-550C	50,000	50 - 150	52 - 70	
44465	TOYOPEARL Hexyl-650C	25	50 - 150	30 - 50	
19026	TOYOPEARL Hexyl-650C	100	50 - 150	30 - 50	
19027	TOYOPEARL Hexyl-650C	1,000	50 - 150	30 - 50	
19028	TOYOPEARL Hexyl-650C	5,000	50 - 150	30 - 50	
21973	TOYOPEARL Hexyl-650C	50,000	50 - 150	30 - 50	



TSKgel HIC resins:

Part #	Product description	Container size (mL)	Bead diameter (µm)	Typical lysozyme capacity (g/L)	
43276	TSKgel Ether-5PW (20)	25	15 - 25	10 - 30	
16052	TSKgel Ether-5PW (20)	250	15 - 25	10 - 30	
16053	TSKgel Ether-5PW (20)	1,000	15 - 25	10 - 30	
18437	TSKgel Ether-5PW (20)	5,000	15 - 25	10 - 30	
43176	TSKgel Ether-5PW (30)	25	20 - 40	10 - 30	
16050	TSKgel Ether-5PW (30)	250	20 - 40	10 - 30	
16051	TSKgel Ether-5PW (30)	1,000	20 - 40	10 - 30	
18439	TSKgel Ether-5PW (30)	5,000	20 - 40	10 - 30	
43277	TSKgel Phenyl-5PW (20)	25	15 - 25	10 - 30	
14718	TSKgel Phenyl-5PW (20)	250	15 - 25	10 - 30	
14719	TSKgel Phenyl-5PW (20)	1,000	15 - 25	10 - 30	
18438	TSKgel Phenyl-5PW (20)	5,000	15 - 25	10 - 30	
43177	TSKgel Phenyl-5PW (30)	25	20 - 40	10 - 30	
14720	TSKgel Phenyl-5PW (30)	250	20 - 40	10 - 30	
14721	TSKgel Phenyl-5PW (30)	1,000	20 - 40	10 - 30	
17210	TSKgel Phenyl-5PW (30)	5,000	20 - 40	10 - 30	

